An algebraic formulation of the graph reconstruction conjecture

Igor C. Oliveira* Columbia University oliveira@cs.columbia.edu Bhalchandra D. Thatte[†] Universidade de São Paulo thatte@ufmg.br

September 9, 2014

Abstract

The graph reconstruction conjecture asserts that every finite simple graph on at least three vertices can be reconstructed up to isomorphism from its deck - the collection of its vertex-deleted subgraphs. Kocay's Lemma is an important tool in graph reconstruction. Roughly speaking, given the deck of a graph G and any finite sequence of graphs, it gives a linear constraint that every reconstruction of G must satisfy.

Let $\psi(n)$ be the number of distinct (mutually non-isomorphic) graphs on n vertices, and let d(n) be the number of distinct decks that can be constructed from these graphs. Then the difference $\psi(n) - d(n)$ measures how many graphs cannot be reconstructed from their decks. In particular, the graph reconstruction conjecture is true for *n*-vertex graphs if and only if $\psi(n) = d(n)$.

We give a framework based on Kocay's lemma to study this discrepancy. We prove that if M is a matrix of covering numbers of graphs by sequences of graphs, then $d(n) \ge \operatorname{rank}_{\mathbb{R}}(M)$. In particular, all *n*-vertex graphs are reconstructible if one such matrix has rank $\psi(n)$. To complement this result, we prove that it is possible to choose a family of sequences of graphs such that the corresponding matrix M of covering numbers satisfies $d(n) = \operatorname{rank}_{\mathbb{R}}(M)$.

1 Introduction

The graph reconstruction conjecture was proposed by Ulam [14] and Kelly [4]. Informally, it states that if two finite, undirected, simple graphs on at least three vertices have the same collection (*multi-set* or *deck*) of *unlabelled* vertex-deleted subgraphs, then the graphs are isomorphic; in other words, any such graph can be *reconstructed* up to isomorphism from the collection of its unlabelled vertex-deleted subgraphs.

The conjecture has been verified by McKay [8] for all undirected, finite, simple graphs on eleven or fewer vertices. In addition, it has been proven for many particular classes of graphs, such as regular graphs, disconnected graphs and trees (Kelly [5]). In fact, Bollobás [2] showed that for almost all graphs, just three (carefully chosen) subgraphs in the deck are sufficient to reconstruct the graph. On the other hand, a similar conjecture does not hold for directed graphs: Stockmeyer

^{*}Supported in part by NSF grants CCF-0915929 and CCF-1115703.

[†]Supported by CNPq grant 151782/2010-5 and by MaCLinC Project at Universidade de São Paulo.

[11, 12] constructed a number of infinite families of non-reconstructible directed graphs. For a more comprehensive introduction to the problem, we refer to a survey by Bondy [3]. For the standard graph theoretic terminology not defined here, we refer to West [15].

Kelly's Lemma [5] is one of the most useful results in graph reconstruction. Let s(F, G) denote the number of subgraphs of G isomorphic to F. Kelly's lemma states that for v(F) < v(G), the parameter s(F, G) is reconstructible, in the sense that if G and G' have the same deck then s(F, G') = s(F, G). Several propositions in graph reconstruction rely on this useful lemma.

Kocay's Lemma [6] allows us, to some extent, to overcome the restriction v(F) < v(G) in Kelly's lemma. It provides a linear constraint on $s(\cdot, G)$ that must be satisfied by every reconstruction of G. Informally, it says that, if $\mathcal{F} = (F_1, \ldots, F_m)$ is a sequence of graphs, each of which has at most v(G) - 1 vertices, then there are constants $c(\mathcal{F}, H)$ such that the value of the sum $\sum_H c(\mathcal{F}, H) \cdot s(H, G)$ is reconstructible, where the sum is taken over all unlabelled *n*-vertex graphs H. Roughly speaking, the constant $c(\mathcal{F}, H)$ counts the number of ways to *cover* the graph H by graphs in the sequence \mathcal{F} .

Kocay's Lemma has been used to show several interesting results in graph reconstruction. For instance, by carefully selecting the sequence \mathcal{F} , it is possible to give a simple proof that disconnected graphs are reconstructible. In addition, it can be used to show that the number of perfect matchings, the number of spanning trees, the characteristic polynomial, the chromatic polynomial, and many other parameters of interest are reconstructible; see Bondy [3].

It is natural to wonder whether even more restrictions may be imposed on the reconstructions of G by applications of Kocay's Lemma. Recall that it is possible to use different sequences of graphs in each invocation of the lemma, and as explained before, for each sequence we get a linear constraint that the reconstructions of G must satisfy. By analysing such equations one would expect to obtain a wealth of information about the structure of any reconstruction of G (perhaps enough equations may even allow us to conclude that G is reconstructible). In this paper we investigate how much information one can obtain by setting up such equations.

We prove that the equations obtained by applying Kocay's Lemma to the deck of a graph G using distinct sequences of graphs provide important information not only about the reconstructions of G, but also on the total number of non-reconstructible graphs on n vertices. More formally, let d(n)be the number of distinct decks obtained from n-vertex graphs. We show that if M is the matrix of coefficients corresponding to these equations, then $d(n) \geq \operatorname{rank}_{\mathbb{R}}(M)$, i.e., the rank of this matrix provides a lower bound on the number of distinct decks. In particular, the existence of a full-rank matrix of coefficients would imply that all graphs on n vertices are reconstructible. In addition, we give a proof that there exist d(n) sequences of graphs $\mathcal{F}_1, \mathcal{F}_2, \ldots, \mathcal{F}_{d(n)}$, with corresponding matrix M of covering numbers, such that $\operatorname{rank}_{\mathbb{R}}(M) = d(n)$. In other words, if the graph reconstruction conjecture holds for graphs with n vertices, then there is a corresponding full-rank matrix certifying this statement.

We state our results in more generality for graphs, hypergraphs, directed graphs, and also for classes of graphs for which similar equations can be constructed; for example, analogous results hold for planar graphs, disconnected graphs and trees.

Similar system of equations where considered by Kocay [7], where he restricted the total number of edges appearing in each sequence of graphs on a given system of equations to be the same. Interestingly, in this case it is not possible to show an equivalence to the graph reconstruction conjecture. In particular, Kocay computed the ratio of the number of independent edge-identities and the number of mutually non-isomorphic graphs with v vertices, e edges, and no isolated vertices (for small parameters v, e), and observed that these values can be strictly less than one. Kocay asked if the reconstruction conjecture would fail to be true if the ratio became small enough.

Our contribution may be summarised as follows. We remove the restriction, as in Kocay's paper [Theorem 5.2 in 6], that the total number of edges be fixed among the sequences of graphs used to derive edge-identities. We show that the number of independent equations available at our disposal is precisely the number of distinct decks on a given number of vertices. Thus we give an algebraic characterisation, based on Kocay's lemma, for the discrepancy between the number of different decks and the number of distinct graphs - a measure of how badly Ulam's conjecture would fail to hold, if indeed it were to be false. In view of the result of Bollobás mentioned earlier, the ratio of the number of independent equations and the number of distinct graphs cannot be small.

A different mathematical perspective on such equations is presented in Mnukhin [10], where reconstruction problems are discussed in the more general context of orbit algebras. Mnukhin's paper also mentions a formulation of Ulam's conjecture in algebraic terms, based on whether the graph algebra is generated by disconnected graphs only. While there may be a translation between the two formulations, this is not immediately obvious to the authors. We refer the reader to Mnukhin's paper for further details, and to the original reference [9] (in Russian) discussed in [10]. Our results are simple to prove, can be specialised to several classes of graphs, as well as generalised to digraphs and hypergraphs, and provide an exact characterisation of the maximum number of independent equations.

Finally, our results may also be viewed as a limitation of the lemmas of Kelly and Kocay (which is proved using Kelly's lemma), and in this regard we share the pessimism expressed by Tutte (see Chapter 9, page 113, [13]). The fact that the number of independent equations is equal to the number of decks suggests that the difficulties with Ulam's conjecture lie somewhere else. In particular, it seems unlikely that applications of Kelly's lemma and Kocay's lemma will shed light on these difficulties.

2 Preliminaries

In this paper, we consider general finite graphs - undirected graphs, directed graphs, hypergraphs, graphs with or without multiple edges, and with or without loops. We take the vertex set of a graph to be a finite subset of \mathbb{N} . We write $V^{(k)}$ for the family of k-element subsets of a set V. Further, we use the notation $v(G) \stackrel{\text{def}}{=} |V(G)|$ and $e(G) \stackrel{\text{def}}{=} |E(G)|$.

Definition 2.1 (Graphs). A hypergraph G is a triple (V, E, ϕ) , where V is its vertex set (also called ground set, and written as V(G)) and E is its set of hyperedges (written as E(G)), and a map $\phi: E \to 2^V \setminus \emptyset$. An undirected graph G is a hypergraph with the restriction that $\phi: E \to V^{(1)} \cup V^{(2)}$; in this case we call a hyperedge e an edge (if $|\phi(e)| = 2$) or a loop (if $|\phi(e)| = 1$). An undirected graph is simple if it contains no loop. A directed graph G is a triple (V, E, ψ) , where V is its vertex set and E is the set of its arcs, and a map $\psi: E \to V \times V$. The first element of $\psi(e)$ is called the tail of the arc e, and the second element of $\psi(e)$ is called the head of e. We denote the set of all finite graphs (including hypergraphs, undirected graphs and directed graphs) by \mathcal{G}^* .¹

Remark 2.2. Although our results and proofs are stated in full generality, it may be helpful in a first reading to consider only finite, simple, undirected graphs.

¹Observe that we are defining graphs using triples because multiple edges are allowed.

Definition 2.3 (Graph isomorphism). Let G and H be two graphs. We say that G and H are *isomorphic* (written as $G \cong H$) if there are one-one maps $f: V(G) \to V(H)$ and $g: E(G) \to E(H)$ such that an edge e and a vertex v are incident in G if and only the edge g(e) and the vertex f(v) are incident in H. Additionally, in the case of directed graphs, a vertex v is the head (or the tail) of an arc e if and only if f(v) is the head (or, respectively, the tail) of g(e). The isomorphism class of a graph G, denoted by G/\cong , is the set of graphs isomorphic to G.

Definition 2.4. A *class of graphs* is a set of graphs that is closed under isomorphism. A class of graphs is said to be *finite* if contains finitely many isomorphism classes.

Definition 2.5 (Reconstruction). Let G be graph and let v be a vertex of G. The induced subgraph of G obtained by deleting v and all edges incident with v is called a vertex-deleted subgraph of G, and is written as G - v. We say that H is a reconstruction of G (written as $H \sim G$) if there is a one-one map $f: V(G) \to V(H)$ such that for all $v \in V(G)$, the graphs G - v and H - f(v) are isomorphic. The relation \sim is an equivalence relation. We say that a graph G is reconstructible if every reconstruction of G is isomorphic to G (i.e., if $H \sim G$ implies $H \cong G$). A parameter t(G) is said to be reconstructible if t(H) = t(G) for all reconstructions H of G. Let C be a class of graphs. We say that C is recognisable if, for any $G \in C$, every reconstruction of G is in C. Furthermore, we say that C is reconstructible if every graph $G \in C$ is reconstructible.

Example 2.6. Let $G(V, E, \phi)$ be a hypergraph. The number of edges incident with all vertices (i.e., edges $e \in E$ such that $\phi(e) = V$, which we call *big edges*), is not a reconstructible parameter. For example, if G^k is a graph obtained from G by adding k new edges e_1, e_2, \ldots, e_k and making them incident with all vertices in V, then G^k is a reconstruction of G. In this sense, no hypergraphs are reconstructible, and each hypergraph has infinitely many mutually non-isomorphic reconstructions. If G is a graph in class C, then C is not recognisable if for some k, the graph G^k is not in C; and C is not finite if graphs G^k are all in C. On the other hand, the number of *small edges*, i.e., edges $e \in E$ such that $\phi(e) \neq V$, is a reconstructible parameter.

In view of the above example, we will always use \mathcal{G}^* for the set of all graphs, \mathcal{G} for the set of all graphs without big edges, and \mathcal{G}_n for the set of *n*-vertex graphs without big edges. A class \mathcal{C}_n will always be a subset of \mathcal{G}_n . We will use the following slightly restrictive definitions for some other reconstruction terms.

Definition 2.7. A graph G in \mathcal{G} is reconstructible if it is *reconstructible modulo big edges*, i.e., if G' is a reconstruction of G and $G' \in \mathcal{G}$, then G' is isomorphic to G. A subclass \mathcal{C} of \mathcal{G} is recognisable if for each graph G in \mathcal{C} , each reconstruction of G in \mathcal{G} is also in \mathcal{C} . A subclass \mathcal{C} of \mathcal{G} is reconstructible if each graph in \mathcal{C} is reconstructible (modulo big edges).

Example 2.8. Disconnected undirected graphs on 3 or more vertices are recognisable and reconstructible. However, there are classes of graphs that are recognisable, but not known to be reconstructible. An important example is the class of planar graphs (Bilinski et al. [1]).

Since \cong and \sim are equivalence relations, the *quotient* notation may be conveniently used to define various equivalence classes of graphs. We write the set of all isomorphism classes of graphs as \mathcal{G}^*/\cong ; analogously we use \mathcal{G}_n/\cong , \mathcal{C}/\cong , \mathcal{C}_n/\cong , and so on. We define an *unlabelled graph* to be an isomorphism class of graphs. But sometimes we abuse the notation slightly, e.g., if a quantity is invariant over an isomorphism class H, then in the same context we may also use H to mean

a representative graph in the class. Similarly, we denote various reconstruction classes by \mathcal{G}/\sim , \mathcal{G}_n/\sim , \mathcal{C}_n/\sim , \mathcal{C}_n/\sim , and so on. Note that equivalence classes of any class of graphs under \sim are refined by \cong ; in particular, $|\mathcal{C}_n/\sim| \leq |\mathcal{C}_n/\cong|$, and equality holds if and only if the class \mathcal{C}_n is reconstructible. We will refer to reconstruction classes of \mathcal{C}_n (i.e., members of \mathcal{C}_n/\sim) by R_1, R_2, \ldots , and isomorphism classes of R_i (i.e., members of R_i/\cong) by $R_{i,1}, R_{i,2}, \ldots$.

Given graphs G and H, the number of subgraphs of G isomorphic to H is denoted by s(H,G). The following two subgraph counting lemmas are important results about the reconstructibility of the parameter s(H,G).

Lemma 2.9 (Kelly's Lemma, [5]). Let H be a reconstruction of G. If F is any graph such that v(F) < v(G), then s(F,G) = s(F,H).

Definition 2.10. Let G be a graph and let $\mathcal{F} := (F_1, F_2, \ldots, F_m)$ be a sequence of graphs. A cover of G by \mathcal{F} is a sequence (G_1, G_2, \ldots, G_m) of subgraphs of G such that $G_i \cong F_i$, $1 \le i \le m$, and $\bigcup G_i = G$. The number of covers of G by \mathcal{F} is denoted by $c(\mathcal{F}, G)$.

Lemma 2.11 (Kocay's Lemma, [6]). Let G be a graph on n vertices. For any sequence of graphs $\mathcal{F} := (F_1, F_2, \ldots, F_m)$, where $v(F_i) < n, 1 \le i \le m$, the parameter

$$\sum_{H} c(\mathcal{F}, H) s(H, G)$$

is reconstructible, where the sum is over all unlabelled n-vertex graphs H.

Proof. We count in two ways the number of sequences (G_1, \ldots, G_m) of subgraphs of G such that $G_i \cong F_i, 1 \le i \le m$. We have

$$\prod_{i=1}^{m} s(F_i, G) = \sum_X c(\mathcal{F}, X) s(X, G),$$
(1)

where the sum extends over all unlabelled graphs X on at most n vertices. Since $v(F_i) < n$, it follows by Kelly's Lemma that the left-hand side of this equation is reconstructible. On the other hand, the terms $c(\mathcal{F}, X)s(X, G)$ are also reconstructible whenever v(X) < n. The result follows after rearranging Equation 1.

To state our results in full generality, we make the following definition.

Definition 2.12. Let C_n be a class of graphs on n vertices. We say that C_n satisfies Kocay's lemma if, for every graph $G \in C_n$ and every sequence of graphs $\mathcal{F} = (F_1, F_2, \ldots, F_m)$, where $v(F_i) < n$, $1 \le i \le m$, the sum

$$\sum_{H \in \mathcal{C}_n/\cong} c(\mathcal{F}, H) s(H, G)$$

is reconstructible.

The following proposition gives a simple condition that is sufficient for a class of graphs C_n to satisfy Kocay's lemma.

Proposition 2.13. Let C_n be a class of graphs on n vertices. Suppose that s(H,G) is reconstructible for every $G \in C_n$ and for every n-vertex graph $H \notin C_n$. Then the class C_n satisfies Kocay's lemma.

Proof. Let $G \in C_n$. Let $\mathcal{F} := (F_1, F_2, \ldots, F_m)$ be any sequence of graphs such that $v(F_i) < n$, $1 \le i \le m$. We write the R.H.S. of Equation 1 as

$$\sum_{H \in \mathcal{C}_n/\cong} c(\mathcal{F},H) s(H,G) + \sum_{H \notin \mathcal{C}_n/\cong} c(\mathcal{F},H) s(H,G),$$

where the second summation is reconstructible. Now we rearrange the terms in Equation 1 to obtain $\sum_{H \in C_n/\cong} c(\mathcal{F}, H) s(H, G)$.

The class of connected simple graphs satisfies Kocay's lemma, since if G is any connected graph and H is any disconnected graph, then s(H,G) is reconstructible (see Bondy [3]). Other classes of graphs that satisfy Kocay's lemma include planar graphs, trees and of course the class of all graphs. Our theorems apply to finite and recognisable classes of graphs satisfying Kocay's Lemma. All the above classes of graphs are recognisable as well.

Let $C_n \subseteq G_n$ be a finite, recognisable class of *n*-vertex graphs satisfying Kocay's Lemma. In the rest of this paper, we study equations obtained by applying Kocay's Lemma to C_n . It is useful to view this lemma as follows. Let $\mathcal{F} := (F_1, \ldots, F_m)$, be a sequence of graphs where $v(F_i) < n$ for each $1 \leq i \leq m$. Let $G, G' \in R \in C_n/\sim$, i.e., G' is a reconstruction of G, and since C_n is recognisable, G' is in C_n . Then we have

$$\sum_{H \in \mathcal{C}_n/\cong} c(\mathcal{F}, H) s(H, G') = k_{\mathcal{F}, R}$$

where $k_{\mathcal{F},R}$ is a constant that depends only on the sequence \mathcal{F} and the reconstruction class R, i.e., it is a reconstructible parameter. In this expression, $c(\mathcal{F}, H)$ is constant (i.e., it is independent of the reconstruction class) and s(H, G') depends on the isomorphism class of a particular reconstruction G' of G under consideration. Therefore, each application of Kocay's Lemma provides a linear constraint on s(H, G') that all reconstructions G' of G must satisfy.

This paper is devoted to a study of systems of such linear constraints obtained by applications of Kocay's lemma. In particular, we study the rank of a matrix of covering numbers that we define next.

Definition 2.14. Let C_n be a finite class of graphs on n vertices. Let $\mathfrak{F} = (\mathcal{F}_1, \mathcal{F}_2, \ldots, \mathcal{F}_l)$ be a family of sequences of graphs on at most n-1 vertices. We let $M_{\mathfrak{F}, C_n/\cong} \in \mathbb{R}^{|\mathfrak{F}| \times |C_n/\cong|}$ to be a matrix whose rows are indexed by the sequences $\mathcal{F}_i, i = 1, 2, \ldots, l$ and whose columns indexed by the distinct isomorphism classes of graphs in C_n . The entries of $M_{\mathfrak{F}, C_n/\cong}$ are the covering numbers defined by $c(\mathcal{F}, H)$, where $\mathcal{F} \in \mathfrak{F}$ and $H \in C_n/\cong$.

3 On the rank of a matrix obtained from Kocay's Lemma

3.1 Large rank implies few non-reconstructible graphs

As observed earlier, for any finite class C_n of graphs, $|C_n/\sim| \leq |C_n/\cong|$, and the bigger the number of distinct reconstruction classes, the smaller is the number of non-reconstructible graphs. The main result of this section, Theorem 3.2, states that for any finite, recognisable class of graphs satisfying Kocay's lemma, the number of distinct reconstruction classes is bounded from below by the rank of the matrix of covering numbers, for any family of sequences of graphs.

Let \mathcal{C}_n be a finite, recognisable class of *n*-vertex graphs satisfying Kocay's Lemma. Let \mathfrak{F} be a finite family of sequences of graphs on at most n-1 vertices. Let $M_{\mathfrak{F},\mathcal{C}_n/\cong}$ be the corresponding matrix of covering numbers $c(\mathcal{F},H)$, where $\mathcal{F} \in \mathfrak{F}$ and $H \in \mathcal{C}_n/\cong$ (see Definition 2.14). Let $W = \{x \in \mathbb{R}^{|\mathcal{C}_n/\cong|} \mid M_{\mathfrak{F},\mathcal{C}_n/\cong} \cdot x \equiv 0\}$ be a subspace of the vector space $\mathbb{R}^{|\mathcal{C}_n/\cong|}$ over \mathbb{R} . We associate with \mathcal{C}_n the constant $\alpha(\mathcal{C}_n) \coloneqq |\mathcal{C}_n/\cong| - |\mathcal{C}_n/\sim|$.

Lemma 3.1. dim
$$(W) \ge \alpha(\mathcal{C}_n)$$
.

Proof. If $\alpha(\mathcal{C}_n) = 0$, the result is trivial. Otherwise, let $R_1, \ldots, R_s \in \mathcal{C}_n/\sim$ be the non-reconstructible reconstruction classes in \mathcal{C}_n , i.e., $r_i \coloneqq |R_i/\cong| > 1$ for all $i \in \{1, 2, \ldots, s\}$. Let $R_{i,j}, j \in \{1, 2, \ldots, r_i\}$ be the isomorphism classes in $R_i, i \in \{1, 2, \ldots, s\}$. Let $G_{i,j}$ be representative graphs from $R_{i,j}$.

For each $G_{i,j}$, we define a vector $w^{i,j} \in \mathbb{R}^{|\mathcal{C}_n/\cong|}$, with its entries, which are indexed by unlabelled graphs $H \in \mathcal{C}_n/\cong$, defined as follows:

$$w^{i,j}(H) := s(H, G_{i,j}) - s(H, G_{i,1}), \text{ where } H \in \mathcal{C}_n / \cong \mathcal{C}_n$$

Observe that to prove the lemma it is enough to show that the vectors $w^{i,j}$ satisfy the following properties:

- (i) for all $i \in \{1, 2, \dots, s\}$, for all $j \in \{1, 2, \dots, r_i\}$, $w^{i,j} \in W$; and
- (*ii*) the vectors in the set $U \coloneqq \{w^{i,j} \mid 1 \leq i \leq s, 2 \leq j \leq r_i\}$ are non-zero and linearly independent, where $|U| = \alpha(\mathcal{C}_n)$.

Proof of (i): Graphs $G_{i,j}$ and $G_{i,1}$ are reconstructions of each other, and \mathcal{C}_n satisfies Kocay's Lemma. Therefore, for every row $M_{\mathcal{F}}$ of $M_{\mathfrak{F},\mathcal{C}_n/\cong}$, we have,

$$\sum_{H \in \mathcal{C}_n/\cong} c(\mathcal{F}, H) s(H, G_{i,j}) = \sum_{H \in \mathcal{C}_n/\cong} c(\mathcal{F}, H) s(H, G_{i,1})$$

$$\therefore \quad M_{\mathcal{F}} \cdot w^{i,j} = \sum_{H \in \mathcal{C}_n/\cong} c(\mathcal{F}, H) (s(H, G_{i,j}) - s(H, G_{i,1})) = 0.$$

Therefore, $M_{\mathfrak{F}, \mathcal{C}_n/\cong} \cdot w^{i,j} = 0.$

Proof of (ii): Let the vectors in U be ordered $u^1, u^2, \ldots, u^{\alpha(\mathcal{C}_n)}$ so that the corresponding graphs are ordered by non-decreasing numbers of small edges. We prove that u^1 is non-zero, and for each $k \in \{2, \ldots, \alpha(\mathcal{C}_n)\}$, the vector u^k is non-zero and is linearly independent of $u^1, u^2, \ldots, u^{k-1}$, which would imply that the vectors in U are linearly independent.

Let $u^{\ell} = w^{i,j}$ for some $i \in \{1, 2, ..., s\}$ and $j \in \{2, ..., r_i\}$. First recall that C_n is recognisable, $R_i \in C_n/\sim$, and $G_{i,j} \in R_i/\cong$; therefore, $G_{i,j} \in C_n/\cong$. In addition, $G_{i,j} \not\cong G_{i,1}$ since $j \ge 2$ and these two graphs belong to distinct isomorphism classes within the same reconstruction class R_i . Finally, the number of small edges is reconstructible, i.e., $e(G_{i,j}) = e(G_{i,1})$. Therefore,

$$u^{\ell}(G_{i,j}) = w^{i,j}(G_{i,j}) = s(G_{i,j}, G_{i,j}) - s(G_{i,j}, G_{i,1}) = 1 - 0 = 1.$$

Now consider the vectors $u^k = w^{i',j'}$ and $u^\ell = w^{i,j}$, where $1 \le k < \ell$. We prove that $u^k(G_{i,j}) = 0$. Since $k < \ell$, according to the ordering of U, we have $e(G_{i',j'}) \le e(G_{i,j})$. Since $G_{i',j'}$ and $G_{i',1}$ are reconstructions of each other, we have $e(G_{i',j'}) = e(G_{i',1})$. Now, if $e(G_{i',j'}) < e(G_{i,j})$, then

$$u^{k}(G_{i,j}) = w^{i',j'}(G_{i,j}) = s(G_{i,j}, G_{i',j'}) - s(G_{i,j}, G_{i',1}) = 0 - 0 = 0.$$

On the other hand, if $e(G_{i',j'}) = e(G_{i,j})$, then again $s(G_{i,j}, G_{i',j'}) = 0$ (since $G_{i,j}$ and $G_{i',j'}$ are non-isomorphic but have the same number of edges) and $s(G_{i,j}, G_{i',1}) = 0$ (because j > 1, so $G_{i,j}$ and $G_{i',1}$ are non-isomorphic but have the same number of edges).

Now the lemma follows from $\alpha(\mathcal{C}_n) := |\mathcal{C}_n/\cong| - |\mathcal{C}_n/\sim| = \sum_{i=1}^{s} (r_i - 1) = |U|.$

Theorem 3.2. Let C_n be a finite, recognisable class of n-vertex graphs satisfying Kocay's Lemma. Let \mathfrak{F} be a family of sequences of graphs on at most n-1 vertices. If $M_{\mathfrak{F},C_n/\cong}$ is the corresponding matrix of covering numbers associated with \mathfrak{F} and C_n , then $|\mathcal{C}_n/\sim| \geq \operatorname{rank}_{\mathbb{R}}(M_{\mathfrak{F},C_n/\cong})$.

Proof. Applying the Rank-Nullity Theorem, we have

$$\dim(W) + \operatorname{rank}_{\mathbb{R}}(M_{\mathfrak{F}, \mathcal{C}_n/\cong}) = |\mathcal{C}_n/\cong|.$$

It follows from Lemma 3.1 that

$$\alpha(\mathcal{C}_n) + \operatorname{rank}_{\mathbb{R}}(M_{\mathfrak{F}, \mathcal{C}_n/\cong}) \leq \dim(W) + \operatorname{rank}_{\mathbb{R}}(M_{\mathfrak{F}, \mathcal{C}_n/\cong}) = |\mathcal{C}_n/\cong| \, .$$

Now recalling the definition of $\alpha(\mathcal{C}_n)$, we have

$$|\mathcal{C}_n/\cong| - |\mathcal{C}_n/\sim| + \operatorname{rank}_{\mathbb{R}}(M_{\mathfrak{F},\mathcal{C}_n/\cong}) \le |\mathcal{C}_n/\cong|,$$

which implies that $|\mathcal{C}_n/\sim| \geq \operatorname{rank}_{\mathbb{R}}(M)_{\mathfrak{F},\mathcal{C}_n/\cong}$.

Corollary 3.3. Under the hypotheses of Theorem 3.2, if $\operatorname{rank}_{\mathbb{R}}(M_{\mathfrak{F},\mathcal{C}_n}) = |\mathcal{C}_n/\cong|$ then every graph in \mathcal{C}_n is reconstructible.

Figure 1 illustrates an application of Corollary 3.3 to the class of connected graphs on four vertices. We show six sequences of graphs (indexing rows) and the corresponding covering numbers for each of the six connected graphs on four vertices (indexing the columns). A zero in *i*-th row and the *j*-th column (e.g., most entries in the upper triangle) indicates that there is no way to cover the corresponding graph (indexing a column) by graphs in the corresponding sequence (indexing the row). The matrix has full rank, implying that connected graphs on four vertices are reconstructible.

3.2 The existence of matrices with optimal rank

Theorem 3.4. Let C_n be a recognizable class of n-vertex graphs satisfying Kocay's lemma. Then there exists a family \mathfrak{F} of sequences of graphs with corresponding matrix of covering numbers $M_{\mathfrak{F},C_n/\cong}$ such that $\operatorname{rank}_{\mathbb{R}}(M_{\mathfrak{F},C_n/\cong}) = |\mathcal{C}_n/\sim|$.

Proof. Let \mathfrak{F} be the family of all *inequivalent* sequences of length at most n of (n-1)-vertex graphs. Here we consider two sequences \mathcal{F}_i and \mathcal{F}_j to be inequivalent if for each bijection f from \mathcal{F}_i to \mathcal{F}_j , there is at least one graph F in \mathcal{F}_i for which f(F) is not isomorphic to F. Since the covering numbers for sequences of length 1 in \mathfrak{F} are all 0, we assume that \mathfrak{F} contains only sequences of length at least 2. Let $M_{\mathfrak{F}, \mathcal{C}_n/\cong}$ be the corresponding matrix of covering numbers. We show below that this choice for the family of sequences and its corresponding matrix of covering numbers satisfy the desired property.

	G_1			G_4		G_6
$\mathcal{F}_1 = \left(\begin{array}{c} & & \\ & & \\ & & \\ & & \end{array} \right)$	2	3	0	0	0	0
$\mathcal{F}_2=\left(egin{array}{cccc} egin{array}{ccccc} egin{array}{cccccccc} egin{array}{cccccccccccccccccccccccccccccccccccc$	6	6	0	0	0	0
$\mathcal{F}_3 = \left(\begin{array}{c} \bullet \\ \bullet \end{array} \right)$	0	0	1	0	0	0
$\mathcal{F}_4 = \left(\begin{array}{ccccccc} & & & \\ & & & \\ & & & \\ \end{array} \right) , \left(\begin{array}{cccccccccccccccccccccccccccccccccccc$	36	36	24	24	0	0
$\mathcal{F}_5 = \left(\left[\begin{array}{c} \\ \\ \\ \end{array} \right], \left[\begin{array}{c} \\ \\ \\ \end{array} \right] \right)$	150	150	240	240	120	0
$\mathcal{F}_6 = \left(\begin{array}{c} \bullet \\ \bullet \\ \bullet \end{array} \right), \bullet \\ \bullet$	540	540	1536	1536	1800	720

Figure 1: A full-rank matrix M of covering numbers $c(\mathcal{F}_i, G_j)$ providing a proof through Corollary 3.3 that all connected graphs on four vertices are reconstructible.

For a sequence \mathcal{F} and a graph G, let $c^*(\mathcal{F}, G)$ denote the number of tuples (G_1, G_2, \ldots, G_m) of subgraphs of G with distinct vertex sets such that $G_i \cong F_i$, $1 \le i \le m$, and $\bigcup G_i = G$. We call such covers *non-overlapping*. Correspondingly, we have the matrix $M^*_{\mathfrak{F}, C_n/\cong}$ of non-overlapping covering numbers.

Now let $\mathcal{F} := (F_1, F_2, \dots, F_\ell)$ be a sequence in \mathfrak{F} . We have the following recurrence for $c(\mathcal{F}, G)$:

$$c(\mathcal{F},G) = \sum_{k=2}^{\ell} \sum_{P \in \mathcal{P}_{\ell}^{k}} \sum_{\mathcal{H}:=(H_{1},H_{2},\dots,H_{k})} \gamma(\mathcal{H})c^{*}(\mathcal{H},G) \prod_{i=1}^{k} c(\mathcal{F}|_{P^{-1}(i)},H_{i}),$$

where \mathcal{P}_{ℓ}^k denotes the set of all onto functions from $\{1, 2, \ldots, \ell\}$ to $\{1, 2, \ldots, k\}$, and $\mathcal{F}|_{P^{-1}(i)}$ is the subsequence of \mathcal{F} consisting of $F_j; j \in P^{-1}(i)$, and the innermost sum is over all inequivalent sequences \mathcal{H} of length k of graphs on (n-1) vertices. This may be explained as follows. Each cover $(G_1, G_2, \ldots, G_{\ell})$ of G by \mathcal{F} naturally corresponds to a partition of $\{1, 2, \ldots, \ell\}$ in k blocks for some $k \in [2..\ell]$, so that i, j are in the same partition if and only if graphs G_i and G_j have the same vertex set. We denote partitions of $\{1, 2, \ldots, \ell\}$ in k blocks by onto maps P from $\{1, 2, \ldots, \ell\}$ to $\{1, 2, \ldots, k\}$ so that the inverse image $P^{-1}(i)$ denotes the *i*-th block. For the *i*-th block $P^{-1}(i)$ of an onto map P, the union of graphs $G_j; j \in P^{-1}(i)$ is a graph H_i on n-1 vertices. We denote the subsequence of \mathcal{F} with indices $j \in P^{-1}(i)$ by $\mathcal{F}|_{P^{-1}(i)}$. Now the cover of G by the sequence $\mathcal{H} := (H_1, H_2, \ldots, H_k)$ is non-overlapping, and each H_i may be covered by $F_j; j \in P^{-1}(i)$ in $c(\mathcal{F}|_{P^{-1}(i)}, H_i)$ ways. We do not need to consider the trivial partition of $\{1, 2, \ldots, \ell\}$ into a single block, because there is no cover $(G_1, G_2, \ldots, G_\ell)$ of G by \mathcal{F} such that all G_i have the same vertex set. In other words, the above formula computes $c(\mathcal{F}, G)$ by partitioning the coverings according to k, P, and \mathcal{H} , and then counting the number of coverings in each block of the partition. Since in the formula we use onto functions instead of partitions, the same block of coverings under this partition may be counted more than once, and therefore there is factor $\gamma(\mathcal{H})$ in the formula. If sequence \mathcal{H} contains k_1 copies of a graph Γ_1 , k_2 copies of a graph Γ_2 , and so on, where Γ_i are mutually non-isomorphic graphs, then $\gamma(\mathcal{H}) = (\prod_i k_i!)^{-1}$.

Now we rearrange the terms and write

$$c^{*}(\mathcal{F},G) = c(\mathcal{F},G) - \sum_{k=2}^{\ell-1} \sum_{P \in \mathcal{P}_{\ell}^{k}} \sum_{\mathcal{H}:=(H_{1},H_{2},\dots,H_{k})} \gamma(\mathcal{H})c^{*}(\mathcal{H},G) \prod_{i=1}^{k} c(\mathcal{F}|_{P^{-1}(i)},H_{i})$$

Thus we have expressed the non-overlapping covering numbers for a sequence of length ℓ of graphs in terms of the non-overlapping covering numbers for sequences of length at most $\ell-1$. In the above equation, $c(\mathcal{F}|_{P^{-1}(i)}, H_i)$ are constants independent of G. Also, if $\ell = 2$, we have $c^*(\mathcal{F}, G) = c(\mathcal{F}, G)$. Therefore, by repeatedly applying the above equation to terms containing non-overlapping covering numbers, we eventually obtain

$$c^*(\mathcal{F}, G) = \sum_{\mathcal{F}'} \beta_{\mathcal{F}}(\mathcal{F}') c(\mathcal{F}', G).$$

We have written the coefficients as $\beta_{\mathcal{F}}(\mathcal{F}')$ to emphasize that they arise from factors $c(\mathcal{F}|_{P^{-1}(i)}, H_i)$ and $\gamma(\mathcal{H})$ that do not depend on G. That is, the linear dependence of the non-overlapping covering numbers on the covering numbers is the same for all graphs (but of course depends on \mathcal{F}). Therefore, we can write

$$c^*(\mathcal{F}, \cdot) = \sum_{\mathcal{F}'} \beta_{\mathcal{F}}(\mathcal{F}') c(\mathcal{F}', \cdot).$$

In this manner we have shown that the rows of $M^*_{\mathfrak{F}, \mathcal{C}_n/\cong}$ are in the span of the rows of $M_{\mathfrak{F}, \mathcal{C}_n/\cong}$. Therefore, we have

$$\operatorname{rank}_{\mathbb{R}}(M^*_{\mathfrak{F},\mathcal{C}_n/\cong}) \leq \operatorname{rank}_{\mathbb{R}}(M_{\mathfrak{F},\mathcal{C}_n/\cong}).$$

To show that the rank of $M^*_{\mathfrak{F}, \mathcal{C}_n/\cong}$ is $|\mathcal{C}_n/\sim|$, we construct a square submatrix K of $M^*_{\mathfrak{F}, \mathcal{C}_n/\cong}$ as follows. Let $\{R_i, i = 1, 2, \ldots\} \coloneqq \mathcal{C}_n/\sim$. First, for each reconstruction class $R_i, i = 1, 2, \ldots$, we choose one reconstruction G_i arbitrarily from R_i/\cong . For each $i = 1, 2, \ldots$, we keep the row indexed by the sequence (say \mathcal{F}_i) that is equivalent to the sequence $(G_i - v, v \in V(G_i))$, where the vertices of G_i may be ordered arbitrarily, and we keep the column indexed by G_i . We delete all other rows and columns of $M^*_{\mathfrak{F}, \mathcal{C}_n/\cong}$. We show that K has full rank, which will imply that $\operatorname{rank}_{\mathbb{R}}(M^*_{\mathfrak{F}, \mathcal{C}_n/\cong}) \geq \operatorname{rank}_{\mathbb{R}}(K) = |\mathcal{C}_n/\sim|.$

We define a partial order \leq on C_n/\sim so that $R_i \leq R_j$ if there exists a bijection f from $V(G_i)$ to $V(G_j)$ such that for each v in $V(G_i)$, the graph $G_i - v$ is isomorphic to a subgraph of $G_j - f(v)$.

First we verify that the above relation \leq is a partial order on C_n/\sim . The reflexivity and the transitivity are straightforward to verify. We now verify antisymmetry. Let f be a bijection as in the above paragraph. Therefore, for each $v \in V(G_i)$, we have $e(G_i - v) \leq e(G_j - f(v))$. Let

g be a similar bijection from $V(G_j)$ to $V(G_i)$. Therefore, the bijective composition $g \circ f$ from $V(G_i)$ to $V(G_i)$ is such that for all v in $V(G_i)$, we have $G_i - v$ is isomorphic to a subgraph of $G_i - (g \circ f)(v)$, implying that $e(G_i - v) \leq e(G_j - f(v)) \leq e(G_i - (g \circ f)(v))$. Now observe that $\sum_v e(G_i - v) = \sum_v e(G_i - (g \circ f)(v))$, since $g \circ f$ is a bijection from $V(G_i)$ onto itself. Therefore, we must have $e(G_i - v) = e(G_j - f(v))$ for all $v \in V(G_i)$, implying that $G_i - v$ and $G_j - f(v)$ are isomorphic for all $v \in V(G_i)$. In other words, $R_i = R_j$.

We sort the rows and the columns of K so that if $R_i < R_j$, then G_j is to the right of G_i , and the row corresponding to the sequence \mathcal{F}_i is above the row corresponding to the family \mathcal{F}_j .

Now if $c^*(\mathcal{F}_i, G_j) > 0$ then $R_i < R_j$, therefore, the matrix K is upper-triangular. Also, $c^*(\mathcal{F}_i, G_i) > 0$ for all G_i . Therefore, K has full rank; in fact $\operatorname{rank}_{\mathbb{R}}(K)$ is equal $|\mathcal{C}_n/\sim|$. Since the class \mathcal{C}_n is recognizable and satisfies Kocay's lemma, Theorem 3.2 is applicable. Therefore,

$$|\mathcal{C}_n/\sim| = \mathrm{rank}_{\mathbb{R}}(K) \leq \mathrm{rank}_{\mathbb{R}}(M^*_{\mathfrak{F},\,\mathcal{C}_n/\cong}) \leq \mathrm{rank}_{\mathbb{R}}(M_{\mathfrak{F},\,\mathcal{C}_n/\cong}) \leq |\mathcal{C}_n/\sim|\,,$$

which implies the claim for our choice of \mathfrak{F} , and the corresponding matrix $M_{\mathfrak{F}, \mathcal{C}_n/\cong}$.

Example 3.5. We provide another simple but non-trivial example in directed graphs, which are in general not reconstructible. Figure 2 illustrates a matrix of covering numbers for directed graphs on 3 vertices, with no multi-arcs or loops. Observe that there are 7 distinct graphs in 4 reconstruction classes: G_1 and G_2 are reconstructible; G_3, G_4, G_5 belong to the same reconstruction class; G_6, G_7 belong to the same reconstruction class. The figure shows 4 rows of the matrix corresponding to 4 graph sequences. The rank of the matrix is 4, which is also the number of reconstruction classes. It is possible to verify that the rank cannot be improved by adding more sequences of graphs.




Figure 2: A matrix of covering numbers for directed graphs on 3 vertices. There are 4 reconstruction classes and the rank of the above matrix is also 4.

Acknowledgements

We would like to thank Hiệp Hàn for useful discussions, and the anonymous referees for bringing our attention to the works of Kocay [7] and Mnukhin [10]. The first author is grateful to Yoshiharu Kohayakawa for hosting him at Universidade de São Paulo, and would like to thank Orlando Lee for helpful discussions at an early stage of this work.

References

- M. Bilinski, Y.S. Kwon, and X. Yu. On the reconstruction of planar graphs. Journal of Combinatorial Theory, Series B, 97(5):745–756, 2007.
- B. Bollobás. Almost every graph has reconstruction number three. J. Graph Theory, 14(1): 1–4, 1990. ISSN 0364-9024.
- [3] J.A. Bondy. A graph reconstructor's manual. In Surveys in combinatorics, 1991 (Guildford, 1991), volume 166 of London Math. Soc. Lecture Note Ser., pages 221–252. Cambridge Univ. Press, Cambridge, 1991.
- [4] P.J. Kelly. On isometric transformations. PhD thesis, University of Wisconsin–Madison, 1942.
- [5] P.J. Kelly. A congruence theorem for trees. Pacific Journal of Mathematics, 7(1):961–968, 1957.
- [6] W.L. Kocay. An extension of Kelly's lemma to spanning subgraphs. In Proceedings of the Tenth Manitoba Conference on Numerical Mathematics and Computing, Vol. II (Winnipeg, Man., 1980), volume 31, pages 109–120, 1981.
- [7] W.L. Kocay. Some new methods in reconstruction theory. In *Combinatorial Mathematics IX*, pages 89–114. Springer, 1982.
- [8] B.D. McKay. Small graphs are reconstructible. Australasian Journal of Combinatorics, 15: 123–126, 1997.
- [9] V.B. Mnukhin. On reconstruction of graph polynomials. In 27 Int. Wiss. Kolloq., Ilmenau, 25-29 Okt, pages 87–90, 1982.
- [10] V.B. Mnukhin. The k-orbit reconstruction and the orbit algebra. Acta Applicandae Mathematica, 29(1-2):83–117, 1992.
- [11] P.K. Stockmeyer. The falsity of the reconstruction conjecture for tournaments. J. Graph Theory, 1(1):19–25, 1977. ISSN 0364-9024.
- [12] P.K. Stockmeyer. A census of nonreconstructible digraphs. I. Six related families. J. Combin. Theory Ser. B, 31(2):232–239, 1981. ISSN 0095-8956.
- [13] W.T. Tutte. Graph theory as I have known it. Oxford University Press, 1998.
- [14] S.M. Ulam. A collection of mathematical problems. Interscience Tracts in Pure and Applied Mathematics, no. 8. Interscience Publishers, New York-London, 1960.
- [15] D.B. West. Introduction to graph theory. Prentice Hall, 2001.